Management Guide N



COMMERCIAL LAYERS

Alternative System

NOVOGen **Brown**

WWW.NOVOGEN-LAYERS.COM

TABLE OF CONTENTS

TABLE OF CONTENTS

1.	HO	USING SYSTEMS	1
	1.1.	ALTERNATIVE SYSTEM DESCRIPTION	1
	1.2.	CHOICE OF REARING SYSTEM AND EQUIPMENT	1
2.	FLC	OCK MANAGEMENT DURING THE REARING PERIOD	2
	2.1.	STOCKING DENSITY, DRINKER SPACE AND FEEDING SPACE IN REARING	2
	2.2.	STARTING UP STRATEGY	3
	2.3.	CLIMATE STRATEGY	3
	2.4.	SYSTEM TRAINING IN AVIARY SYSTEM	6
	2.5.	LIGHT TRAINING STRATEGY	8
	2.6.	FEEDING TRAINING STRATEGY	8
	2.7.	WATER TRAINING STRATEGY	9
	2.8.	GRIT AND GRAIN	9
3.	LIG	HTING PROGRAMME	. 10
	3.1.	GENERAL RULES	. 10
	3.2.	VARIOUS HOUSING AND LIGHTING SITUATIONS TO CONSIDER - EXAMPLE PROGRAMME	S12
	3.3.	LIGHTING PROGRAMME IN LIGHT-CONTROLLED REARING HOUSES (<0.5 LUX)	. 14
	3.4.	LIGHTING PROGRAMME IN SEMI – DARK OR OPEN REARING HOUSES (>0.5 LUX)	. 15
	3.5.	LIGHTING PROGRAMME IN HOT CLIMATE (BETWEEN 20° NORTH AND 20° SOUTH)	. 16
4.	FLC	OCK MANAGEMENT DURING THE PRODUCTION PERIOD	. 17
	4.1.	STOCKING DENSITY, DRINKER SPACE AND FEEDING SPACE IN PRODUCTION	. 17
	4.2.	TRANSFER	. 17
	4.3.	LIGHTING PROGRAMME DURING THE PRODUCTION PERIOD	. 17
	4.4.	LIGHT INTENSITY MANAGEMENT DURING THE PRODUCTION PERIOD	. 18
	4.5.	LIGHT TRAINING STRATEGY	. 18
	4.6.	FLOOR EGGS PREVENTION	. 19
	4.7.	MANAGEMENT OF EGG WEIGHT	. 20
	4.8.	MANAGEMENT OF EGG SHELL QUALITY	.21
5.	MO	NITORING BODYWEIGHT AND UNIFORMITY	. 22
	5.1.	BODYWEIGHT CONTROL	. 22
	5.2.	UNIFORMITY CONTROL	. 22
6.	BEA	AK TREATMENT	. 23
	6.1.	BEAK TREATMENT MONITORING	.23
	6.2.	FULLY BEAKED FLOCKS MANAGEMENT	. 24
7.	WA	TER MANAGEMENT	. 25
	7.1.	WATER MANAGEMENT DURING THE REARING PERIOD	.25
	7.2.	WATER QUANTITY	. 25



Commercial layers – Alternative System Management Guide – NOVOgen BROWN Light

TABLE OF CONTENTS

	7.3.	WATER QUALITY	.25
8.	NUT	RITION	. 28
	8.1.	EXAMPLE OF DIET SPECIFICATIONS FOR REARING PERIOD	. 29
	8.2.	EXAMPLE OF DIET SPECIFICATIONS FROM 2% PRODUCTION TO 28 WEEKS	. 30
	8.3.	EXAMPLE OF DIET SPECIFICATIONS FROM 28 TO 45 WEEKS	.31
	8.4.	EXAMPLE OF DIET SPECIFICATIONS FROM 45 TO 70 WEEKS	. 32
	8.5.	EXAMPLE OF DIET SPECIFICATIONS FROM 70 WEEKS TO DEPLETION	. 33
9.	HEA	ALTH PROGRAMME	. 34
	9.1.	VACCINATION	. 34
	9.2.	PARASITES AND INSECTS MONITORING	. 38
10	LITT	ER	.40
	10.1.	LITTER QUALITY	.40
	10.2.	LITTER MANAGEMENT	.40
11	FIN	E DUST MANAGEMENT	.41
12	PAS	TURE MANAGEMENT	.42
	12.1.	OPTIMIZATION OF PASTURE EXPLORATION	.42
	12.2.	MANAGEMENT OF PARASITIC RISK	.43
13	GEN	NERAL FARM RULES	.44
14	CLE	ANING AND DISINFECTION OF POULTRY HOUSES	.45
	14.1.	OPERATIONS PRIOR TO CLEANING	.45
	14.2.	WASHING	.45
	14.3.	PLACING EQUIPMENT BACK INTO THE HOUSE	.46
	14.4.	DISINFECTION	.46
	14.5.	SANITARY PRECAUTIONS	.46
	14.6.	ASSESSING DISINFECTION EFFECTIVENESS	.46
	14.7.	RODENT CONTROL	.46
	14.8.	RESTING PERIOD	.47
	14.9.	BEFORE THE NEW FLOCK ARRIVES	.47

The performance data contained in this document was obtained from results and experience from our own research flocks and flocks of our customers. In no way does the data contained in this document constitute a warranty or guarantee of the same performance under different conditions of nutrition, density or physical or biological environment. In particular (but without limitation of the foregoing) we do not grant any warranties regarding the fitness for purpose, performance, use, nature or quality of the flocks. NOVOGEN makes no representation as to the accuracy or completeness of the information contained in this document.



HOUSING SYSTEMS

1. HOUSING SYSTEMS

1.1. ALTERNATIVE SYSTEM DESCRIPTION

• Floor system:

Birds are able to move freely in the house. They have access to a litter area where they can practice their natural behavior like scratching and dust bathing. The ground surface can totally be on litter or separated between a litter and slatted area. Perches can be provided from an early age.

• Aviary system:

The system provide living space at different levels, which increases the total useable surface. Thus, more birds can be placed per m^2 of floor area than in floor system. The hens can freely move at floor level and vertically in the system. They can use perches and have access to feeders, drinking system, nests on different levels. The floor is covered by litter where birds can perform natural behavior.

Free-Range:

Free range is either a floor or aviary system where birds have access to an outside area.

1.2. CHOICE OF REARING SYSTEM AND EQUIPMENT

The rearing system and equipment have to be carefully chosen according to the equipment used in the production farm.

Feeding and drinking equipment during the rearing period should be as similar as possible to the ones used in the future production system. It will facilitate the transfer and adaptation to the new production house.

For aviary systems, young pullets have to be trained for a quick adaptation after transfer to use the equipment on different levels. It is important that pullets learn to jump to or from the perches and plateforms as well as to sleep in the system during the night.

REARING SYSTEM	RECOMMENDED PRODUCTION SYSTEM
Floor	Floor / Free range
Aviary	Floor / Aviary / Free range



FLOCK MANAGEMENT DURING THE REARING PERIOD 2. FLOCK MANAGEMENT DURING THE REARING PERIOD

Rearing is a key period for successful laying performances. During this period it is necessary to reach the target bodyweight and flock uniformity to prepare the birds for the production period. In addition, pullet must be trained to optimize adaptation to production system. A special care has also to be given to the development of the digestive tract in order to prepare for the fast increase in consumption at the beginning of the production period. A well-managed rearing period has positive effects on:

- Quick adaptation to production system (onset of lay, floor eggs prevention)
- Egg production (peak of lay and persistency)
- Egg quality (egg weight, egg uniformity, shell strength)
- Liveability

To achieve those objectives, it is necessary to respect the following basics:

- Management (stocking density, drinker and feeder space, water, beak trimming...)
- Climate conditions
- Pullet training
- Lighting programme
- Monitoring bodyweight and uniformity
- Nutrition
- Biosecurity and vaccination

2.1. STOCKING DENSITY, DRINKER SPACE AND FEEDING SPACE IN REARING

	From day old aç	to 2 weeks of ge	From 2 to 5 weeks of age		From 6 weeks to transfer	
	Temperate climate	Hot climate	Temperate climate	Hot climate	Temperate climate	Hot climate
Stocking density (1) / useable area (2)	30 birds/m ²	25 birds/m ²	15 birds/m ²	12-15 birds/m ²	12-14 birds/m ²	8-10 birds/m²
Starter drinkers	1 / 80 chicks	1 / 70 chicks	-	-	-	-
Bell drinkers	1 / 150 birds	1 / 150 birds	1 / 100 chicks	1 / 75 chicks	1 / 100 birds	1 / 75 birds
Nipple drinkers	1 / 12 birds	1 / 10 birds	1 / 12 birds	1 / 10 birds	1 / 12 birds	1 / 10 birds
Starting feed pans	1 / 50 chicks		-		-	
Linear chain feeders	2.5 cm	/ birds	4 cm /	birds	6 cm / bird	
Pan feeders	1 / 30 birds		1 / 25 birds		1 / 25 birds	

(1) In Aviary system, stocking density doesn't depend on the age but the moment when the system is opened.

(2) Useable area includes floor, system and platforms areas.



2.2. STARTING UP STRATEGY

ALL SYSTEM

- Before chicks arrive, feed should be accessible to the chicks. Feeders should be well filled and feed (around 100 gr per chick) should be spread over chick paper especially close to the drinking lines. This stimulates the chicks to use the drinking equipment.
- Feed must be regularly renewed to keep it fresh and attractive.
- Before arrival the drinking lines must be checked, so that fresh water is provided and leakage is prevented.
- Triggering the nipples or water cups encourages the birds to drink
- Unload the chicks close to drinkers and feeders.

FLOOR SYSTEM	AVIARY SYSTEM		
 Additional starter drinkers and feeders can be used in the first 2 weeks If brooding takes place in only part of the house, do not exceed a stocking density of 25 to 30 chicks per available m2. Thus allowing chicks to spread quickly over the whole house within the first 7 days. In case of use of circular brooder guards (rings or surrounds): Choose a diameter of 3 to 4 m at day-old but ensure the ring can be enlarged 48 hours after the arrival Ensure the surround can be easily removed after the birds have familiarised themselves with the location of the drinker and feeder systems. Usually, guards can be removed 5 to 7 days after arrival. 	 Chick paper should be placed in the chicks unloading tier (aviary multi-tier system) or on the bottom of the system. Paper should be used until at least 7-10 days of age, it's important to use a good quality paper. In aviary multi-tier system and in the case of coccidiosis vaccination, papers should be kept until at least 3 weeks with regards to re-contamination. Photo: placement of chick paper on arrival 		

2.3. CLIMATE STRATEGY

Before and after arrival and starting up of the chicks, the below information can be used as guidelines for optimal performance. Be aware that those setting depends highly on local climate conditions and must be adapted accordingly.

Before chicks arrival:

- After hatch, a chick is poikilotherm. This means that it can't fully regulate its own bodytemperature. Therefore it is important that the housing climate is supported at such a way that the thermoneutral bodytemperature of 40°C is achieved.
- Raise the house temperature at least 36 hours before chick arrival to 29°C-30°C
- Pre-heat the whole house 30 to 40 hours prior to chick arrival ensuring the floor and system is fully warmed before placement to 35°C.
- Prior to chick arrival ensuring the floor is fully warmed before placement.



- ✤ After chicks arrival:
- Never overheat the chicks and give them a choice within the desired temperature range.
- Take into account the temperature at chick level.
- Depending on the brooder design, place the brooders high enough above the litter (at least 1.5 m) at an angle, to allow for uniform distribution of the chicks.

Check the chicks body temperature after start up to adjust the house temperature by gently touching the chick cloaca with the probe of an ear thermometer. The optimal chick body temperature is 40-41 °C. It is important to measure the body temperature of a sufficient numbers of chicks distributed in different parts of the house.

Photos: Control of chicks temperature





- Check the distribution and behaviour of the chicks to enable you to adapt and manage the temperature of the House:
 - o Good distribution and activity \rightarrow correct temperature and climate
 - o Chicks cuddle and/or avoid large areas of the barn → temperature is too low or presence of air draft
 - o Chicks are panting and are lying on the ground with their wings spread \rightarrow temperature is too high

Scheme: Chicks distribution example





Table: Temperature and relative humidity recommendations

	Under the brooder	Near the circular guard	Room temperature	Relative humidity
Before arrival (2-3 days)	30 °C	30 °C	30 °C	55 – 60 %
Befor arrival (1 days)	35 °C	35 °C	35 °C	55 – 60 %
Week 1	35 – 33 °C	32 – 31 °C	30 – 28 °C	55 – 60 %
Week 2	32 °C	30 – 28 °C	28 – 26 °C	55 – 60 %
Week 3	28 °C	28 – 26 °C	26 – 24 °C	55 – 60 %
Week 4			22 – 20 °C	55 – 60 %
Week 5			22 – 20 °C	60 – 65 %
Week 6			22 – 20 °C	60 – 65 %
Week 7			22 – 20 °C	60 – 70 %
Week 8			21 – 19 °C	60 – 70 %
Till transfer			20 – 18 °C	60 – 70 %

Temperature and relative humidity are important climate parameters, but not the only ones. An optimal ventilation in the rearing house is important to obtain the right oxygen level, a uniform air and temperature disitribution and for the removal of dust and ammonia. If mechanical ventilation is used, the ventilation settings are calculated based on:

- The right temperature based on the age of the chicks
- Constant RH (between 50-70%). Too dry air can damage the oral epithelium.
- An O2 level around 20%
 - o CO2 with direct heating < 5000 ppm
 - o CO2 with indirect heating < 3000 ppm
 - o CO < 10 ppm
 - o Ammonia< 10 ppm
 - o Inspirable dast < 3,5 mg/m³

Based on those settings the recommendations for minimum ventilation results in:

- 0,7 m3 per kg housed in animal, when outside temperature colder than 10°C.
- 1,0 m3 per kg housed animal, when outside temperature warmer than 10°C.

2.4. SYSTEM TRAINING IN AVIARY SYSTEM

Besides obtaining a good bodyweight and uniformity, one of the main target of aviary rearing system is achieving a well trained pullet that moves actively through the house and over the system. After transfer in production house, this training enhance the access to the different levels of the system and so optimize the nesting behaviour as well as access to the water and feeding system.

Aviary rearing systems can roughly be divided in 2 categories: multi-tier (left) and plateform (right). These two different systems require different management.

Schema and photo of Aviary rearing multi-tier systems





Schema and photo of Aviary rearing platform systems







Pullet training in aviary system

Multi-tier system	Platform system
• Multi-tier systems can be composed of up to 4 tiers. This system can be supplied with a movable	 The chicks are always starting up in the system on chick paper, with all the plateforms down.
plateform between the tiers. If a plateform is presented, it is most of the time combined with a drinking line on top.	 A water line can be installed above the movable plateforms and an extra feeding chain on the upper fixed plateform level. From the first week onwards
• After arrival at the farm, the chicks are usually placed in the middle or upper 2 tiers where temperature is usually slightly higher. For the farmer	the movable plateforms will be progressively lifted so that the chicks have to jump to get to the drinking system.
it provides also a better sight on the chicks. Chicks remain for approximatively 4 weeks in their closed tier.	 Chicks remain for approximatively 4 weeks in the system. Thereafter some side plateforms are opened so that the chicks have access to the litter
• After 4 weeks and if the chicks are well developed and active, system can be opened. If a movable plateform is used, it will be placed at a low level so	space on the ground floor. It is possible to use stairs to stimulate the young chicks to come back from the litter space on the system.
that the chicks can use it as a stair to access the opened level. Most of the time a perche is also present in front side of the cage. This will also help the chicks to jump out or in the cages.	• Together with opening of the system, some side plateforms are placed at an angle. This stimulates the chicks at the same time to move upwards to the second fixed plateform level.
• From week 4 to 8, the movable plateform can be adjusted to the behaviour of the pullets. If the condition and development of the pullets is in the standard, the plateform should be progressively lifted up to stimulate the jumping behaviour of the	 In the weeks that follow (approx. 6 – 10) the adjustment of the movable plateforms and the extra feeding chain at the second fixed plateform level should stimulate the hens to move over the complete system.
 Farmer must check the system behaviour of the pullets when the light is off. All of them should be back and stay overnight in the system. Pullets remaining on floor during the night will have higher risk of laying on the floor once in the production house 	• Farmer must check the system behaviour of the pullets when the light is off. All of them should be back and stay overnight in the system. Pullets remaining on floor during the night will have higher risk of laying on the floor once in the production house.

Furhermore, there are other tools to stimulate the system training of the hens, which are **block feeding**, water training and light training.

2.5. LIGHT TRAINING STRATEGY

A specific dimming procedure can be used to stimulate the pullets to move up from the litter area to the system before night time. Herebelow is an example of procedure:

- Step 1: Turn-off the outside lights above the litter space 15 minutes before the total light is switched off. Since pullets are attracted by light, they will move from the litter space towards the system. When the pullets are young (4 to 6 weeks of age) and just released from the system, the farmer should, in addition, daily stimulate the pullets physically at the same time by walking around the system. When it is done from a young age, the switch-off of the outside lights will work as a Pavlov effect on the long term. Birds will get used to the signal and will start to come back in the system.
- Step 2: Switch off the light inside the rearing system 5 minutes before the total house is dark. This is only possible when there is a 'third' lighting level above the system.
- Step 3: This third light level can be dimmed as final step. It will stimulate the pullets to move more upwards in the system.

The total dimming time should be as short as possible. Start with a maximum of 15 minutes and adapt it to the behaviour of the birds. If the procedure is too long, there is a risk that pullets move back from the system towards the litter space.



Figure: 3 steps light dimming procedure in rearing

2.6. FEEDING TRAINING STRATEGY

During the three first weeks, chicks shall have a permanent access to the feed in order to favour the start. Afterwards, it is recommended to progressively implement empty feeders once per day in the middle of the day or afternoon. The last feed distribution shall occur 1 hour before light switch off by filling the chain at 100%. The benefits of this strategy are:

- Preventing selective feed intake and thereby stimulating growth, development and uniformity
- Triggering the pullets to come in the system before light switch off.



2.7. WATER TRAINING STRATEGY

Water training is another tool that can be used to stimulate the pullets to move through the system. Water training should only be started when the growth and development of the hens is in the standard. If all parameters are on standard, it can be started at approximatively 8 weeks of age.

The principal of water training is to shut down a drinking line on one level, so that the hens have to move to another level for drinking. Based on the vaccination scheme, behaviour and condition of the hens the training can be extended at a flexible way.

This practice requires to closely observe birds behavior as well as daily follow up the water consumption to make sure it remains constant.

Water training can only be used in an aviary multi-tier system. In an aviary platform system there is always water training due to the fact that the hens have to jump on the plateau to reach the water.

2.8. GRIT AND GRAIN

To maintain an active feeding behaviour and to help the development of the digestive tract and encourage the birds to scratch the litter, it is advised to give grit and grain to the birds from 4 to 5 weeks of age:

- Grit (insoluble stone particles of 2 to 4 mm): 1 to 3 g per week per bird, distributed over 2 or 3 days.
- Grain (broken maize, wheat): 3 g per bird every day, or every other day



3. LIGHTING PROGRAMME

3.1. GENERAL RULES

Sexual maturity and production are largely influenced by the changes in day length to which pullets are exposed. Carefully chosen lighting programmes will help to optimise the performance of commercial layers. Remember that sexual maturity and bodyweight at sexual maturity influence egg production, egg size, liveability, and egg shell quality. It is difficult to advise a universally optimum and perfect lighting programme. The following lighting programmes are examples and have to be considered as a guideline to help formulate a lighting programme adapted to your own situation.

To establish your own lighting programme, it is important to take into account the following factors:

- Your location (changes in light duration (day length) during the year)
- The characteristics of the rearing unit (light-controlled, semi-dark or open house type)
- Season of the year (increasing or decreasing day length)
- Temperature (light duration at the highest temperature)
- Date of the hatch (what is the natural day length at the bodyweight targeted when light stimulation will take place for onset of lay?)
- Growth of the flock
- Past records of performance obtained in this rearing unit
- Avoid any unwanted external lights in the dark house. It can affect the lighting program and efficiency and induce feather pecking.

✤ LIGHTING PROGRAMME DURING THE FIRST WEEKS OF THE REARING PERIOD

In order to encourage skeletal development and growth, a slow step-down lighting programme is advised for all housing conditions.

The decrease in artificial light duration during the day is then adjusted according to the housing type. For an open house system (and above 20° Latitude), determine the natural light day length that the birds will encounter before 16 weeks of age. This will then help determine the maximum day length the flock will be given and help avoid an unwanted early light stimulation before the flock has matured sufficiently. Early light stimulation will be promoted by a natural increase in day length during rearing.

In dark rearing houses (and when allowed by the local regulation), it is possible to use an intermittent lighting programme during the first two weeks of age. It allows synchronisation of the chicks' behaviour for; eating, drinking and resting. It can have a beneficial effect on the weakest chicks which are stimulated by the stronger ones and improves the flock uniformity. After two weeks, switch to a regular step down lighting programme.



Table: Intermittent lighting programme



LIGHTING PROGRAMME FROM 8 WEEKS OF AGE TO TARGETED AGE OF LIGHT STIMULATION.

In order to control sexual maturity and to avoid early sexual maturity at an inadequate and immature bodyweight, it is important to avoid every day any increase in light duration (due to a natural increase in day length) during this period.

According to the season in a dark house system a stable day length can be used between 7 weeks of age and the age when light stimulation is targeted. The light duration during this period can also be adapted according to the growth of the pullets (10, 11 or 12 hours could be used when growth is slow).

In an open house system, the most difficult system for controlling sexual maturity, the natural day length at which the pullets will be exposed to at 16 weeks of age will determine the light duration at the plateform to avoid any increase of light duration before 16 weeks of age.

INCREASING DAY LENGTH TO STIMULATE EGG PRODUCTION

For all conditions, the main indicator to determine the time of light stimulation is the bodyweight.

LIGHTING PROGRAMME DURING PRODUCTION

Never decrease the artificial light duration during the production period as this can lead to an early decline in egg production.



Graph: Example of lighting programme in light controlled rearing and laying houses

✤ LIGHT INTENSITY

A higher light intensity during the brooding period will encourage growth by promoting higher levels of activity of the flock and a higher feed intake. After 2 or 3 weeks and according to the behaviour of the chicks, the light intensity may be reduced to match the field conditions and the light intensity the birds will be exposed to during the production period (degree of darkness of the rearing house and the laying house).



3.2. VARIOUS HOUSING AND LIGHTING SITUATIONS TO CONSIDER – EXAMPLE PROGRAMMES

Light-controlled rearing house to light-controlled laying house:

- Use a slow step down lighting programme until 6 weeks of age.
- A constant 9 hours day length from 7 weeks to light stimulation (12 hours may be used where needed according to growth)
- Increase the light duration by 2 hours when body weight is between 1,260-1,330 g,
- Add 1 hour and/or 30 minutes per week until 15.30 hours or 16 hours total light duration is obtained.

Light-controlled rearing house to open or semi-dark laying house:

- Use a slow step down lighting programme until 6 weeks of age.
- A constant 9-10 hours day length from 7 to 15 weeks of age.
- Increase light duration by 2 hours when body weight is between 1,260-1,330 g.
- Add 1 hour and/or 30 minutes per week until 15.30 hours or 16 hours total light is obtained.
- Light intensity in rearing should be managed to avoid any dramatic and sudden increase in light intensity at transfer time.

Open or semi-dark rearing house to light-controlled laying house:

- Use a slow step down lighting programme until 6 weeks of age.
- A constant 9-10 hours (or natural day length) from 7 to 15 weeks of age.
- Increase light duration by 2 hours at 1,260-1,330 g of bodyweight when there is a decreasing day length.
- Increase light duration by 1 hour at 1,260-1,330 g of bodyweight when there is an increasing day length.
- Add 1 hour and/or 30 minutes per week until 15.30 hours or 16 hours total light is obtained.
- Light intensity from transfer time should be managed to avoid any dramatic and sudden decrease of light intensity.

Open or semi-dark rearing house to open or semi-dark laying house:

- Use a slow step down lighting programme until 6 weeks of age.
- A constant 9-10 hours (or NDL) day length from 7 to 15 weeks of age.
- Increase light duration by 2 hours at 1,260-1,330 g of bodyweight when there is a decreasing day length.
- Increase light duration by 1 hour at 1,260-1,330 g of bodyweight when there is an increasing day length.
- Make light stimulation more effective by adding the additional hours of light in the morning instead of the evening.
- Add 1 hour and/or 30 minutes per week until 15.30 hours or 16 hours total light is obtained.



In a hot climate:

- Use a slow step down lighting programme until 12 weeks of age.
- A constant natural day length from 12 weeks of age to 2-5% of production.
- Increase light duration by 1 hour and/or 30 minutes from 2-5% of production in the morning.
- Add 1 hour and/or 30 minutes per week until 15.30 hours or 16 hours total light is obtained.
- The light on should be adapted to allow the birds to eat during the cooler part of the day.

Flash Feeding:

When allowed by the local regulation, it is possible to use an additional 1.00 to 1.30 hours of light with feed usually 3 hours after the lights going out, to ensure the longest period of dark remains after the lights go back out in order to promote an optimal feed intake during the first weeks of production or to compensate for the adverse effect of high temperature during the summer.

This extra light period may be introduced and removed during the production period at any time after the increase in light duration at the start of lay.

Please contact the NOVOGEN technician in your area for more specific advice.



3.3. LIGHTING PROGRAMME IN LIGHT-CONTROLLED REARING HOUSES (<0.5 LUX)

Age	Age	Bodvweight at start	Average		
(weeks)	(days)	of the week (g)	laying rate of	Light duration in hours	Light intensity
0	0 to 2			22.00 (1)	20-40 lux
1	3 to 7			20.00 (1)	20-30 lux
2	8 to 14			19.00	10-20 lux
3	15 to 21			17.00	5-10 lux
4	21 to 28			15.00	5-10 lux
5	29 to 35			13.00	5-10 lux
6	36 to 42			11.00	5-10 lux
7	43 to 49			10.00	5-10 lux
8	50 to 56			09.00	5-10 lux
9	57 to 63			09.00	5-10 lux
10	64 to 70			09.00	5-10 lux
11	71 to 77			09.00	5-10 lux
12	78 to 84			09.00	5-10 lux
13	85 to 91			09.00	5-10 lux
14	92 to 98			09.00	5-10 lux
15	99 to 105	(4)		09.00 (4)	5-10 lux
16	106 to 112	1260-1330		11.00	5-10 lux
17	113 to 119	1340-1415		12.00	5-15 lux
18	120 to 126	1410-1490	0-2%	13.00	5-15 lux
19	127 to 133	1480-1560	2-25%	14.00 (2)	5-15 lux
20	134 to 140	1550-1630	30-55%	14.30	5-15 lux
21	141 to 147	1610-1700	55-82%	15.00	5-15 lux
22	148 to 154	1655-1750	80-90%	15.30	5-15 lux
23	155 to 161	1690-1780	85-93%	15.30	5-15 lux
24	162 to 168	1710-1805	88-94%	15.30 (3)	5-15 lux
25	169 to 175	1725-1820	90-95%	15.30 (3)	5-15 lux
25 +	176 to 182			15.30 (3)	5-15 lux

(1) - An intermittent lighting programme can be implemented (if allowed by the local regulation)

(2) - From 19 weeks of age, flash feeding could be added (if allowed by the local regulation)

(3) - Could be increased to 16 hours according to feed consumption

(4) - According to the average egg weight requested by the market, it could be possible to light stimulate the pullets one week earlier

(5) - Light intensity and light stimulation should be adapted to local environment and flock condition.



3.4. LIGHTING PROGRAMME IN SEMI – DARK OR OPEN REARING HOUSES (>0.5 LUX)

Age	Age	Bodyweight at start	Average laying	In decreasing day	In increasing day
(week)	(days)	of the week (g)	rate of the week	length in hours	length in hours
0	0 to 2			22.00	22.00
1	3 to 7			20.00	20.00
2	8 to 14			19.00	19.00
3	15 to 21			17.00	17.00
4	21 to 28			15.00	15.00
5	29 to 35			13.00 (or NDL)	13.00 (or NDL)
6	36 to 42			12.00 (or NDL)	1200. (or NDL)
7	43 to 49			10.00 (or NDL)	10.00 (or NDL)
8	50 to 56			10.00 (or NDL)	10.00 (or NDL)
9	57 to 63			10.00 (or NDL)	10.00 (or NDL)
10	64 to 70			10.00 (or NDL)	10.00 (or NDL)
11	71 to 77			10.00 (or NDL)	10.00 (or NDL)
12	78 to 84			10.00 (or NDL)	10.00 (or NDL)
13	85 to 91			10.00 (or NDL)	10.00 (or NDL)
14	92 to 98			10.00 (or NDL)	10.00 (or NDL)
15	99 to 105			10.00 (or NDL)	10.00 (or NDL)
16	106 to 112	1260-1330		+2.00 (or 16.00)	+1.00 (or 16.00)
17	113 to 119	1340-1415		+1.00 (or 16.00)	+1.00 (or 16.00)
18	120 to 126	1410-1490	0-2%	+1.00 (or 16.00)	+1.00 (or 16.00)
19	127 to 133	1480-1560	2-25%	+1.00 (or 16.00) (1)	+1.00 (or 16.00)
20	134 to 140	1550-1630	30-55%	+0.30 (or 16.00)	+0.30 (or 16.00)
21	141 to 147	1610-1700	55-82%	+0.30 (or 16.00)	+0.30 (or 16.00)
22	148 to 154	1655-1750	80-90%	+0.30 (or 16.00)	+0.30 (or 16.00)
23	155 to 161	1690-1780	85-93%	16.00	+0.30 (or 16.00)
24	162 to 168	1710-1805	88-94%	16.00	16.00
25	169 to 175	1725-1820	90-95%	16.00	16.00
25+	176 to 182				

NDL: Natural Day Length

(1) - From 19 weeks of age, flash feeding could be added



3.5. LIGHTING PROGRAMME IN HOT CLIMATE (BETWEEN 20° NORTH AND 20° SOUTH)

Age (week)	Age (days)	Bodyweight at start	Average laying	Light duration
Age (week)	Age (uays)	of the week (g)	rate of the week	in hours
0	0 to 2			22.00
1	3 to 7			20.00
2	8 to 14			19.00
3	15 to 21			18.00
4	21 to 28			17.00
5	29 to 35			16.00
6	36 to 42			15.30
7	43 to 49			15.00
8	50 to 56			14.30
9	57 to 63			14.00
10	64 to 70			13.30
11	71 to 77			13.00
12	78 to 84			12.30
13	85 to 91			12.00
14	92 to 98			12.00 (or NDL)
15	99 to 105			12.00 (or NDL)
16	106 to 112	1260-1330		12.00 (or NDL)
17	113 to 119	1340-1415		12.00 (or NDL)
18	120 to 126	1410-1490	0-2%	12.00 (or NDL)
19	127 to 133	1480-1560	2-25%	+ 1.00 (1)
20	134 to 140	1550-1630	30-55%	+1.00
21	141 to 147	1610-1700	55-82%	+ 1.00
22	148 to 154	1655-1750	80-90%	+ 0.30
23	155 to 161	1690-1780	85-93%	+0.30 (or 16.00)
24	162 to 168	1710-1805	88-94%	16.00
25	169 to 175	1725-1820	90-95%	16.00
25+	176 to 182			

NDL: Natural Day Length

(1) - From 19 weeks of age, flash feeding could be added (if allowed by the local regulation)



FLOCK MANAGEMENT DURING THE PRODUCTION PERIOD 4. FLOCK MANAGEMENT DURING THE PRODUCTION PERIOD

4.1. STOCKING DENSITY, DRINKER SPACE AND FEEDING SPACE IN PRODUCTION

Temperate climate		Hot climate		
Stocking density / useable area (1)	8-9 birds/m ²	5-6 birds/m ²		
Bell drinkers	1 cm / bird	1,5 cm / bird		
Nipple drinkers	1 / 10 birds	1 / 8 birds		
Linear chain feeders	10 cm / bird			
Pan feeders	1 / 20 birds			
Perches	15 cm / bird			
	(> 30 cm between perches and > 20 cm between wall and perch)			
Pon Holes	2m / 1000 birds			
	(higth > 35cm / width > 40 cm)			
Nest	Individual nest = 1 nest / 7 birds			
1001	Group nest = 1m ^{2 /} 120 birds			
Litter area	> 250 c	m ² / bird		

(1) Useable area includes floor, system and platforms areas. Nests area is not included.

4.2. TRANSFER

Transfer is advised around 16 to 17 weeks of age.

- Before the appearance of the 1st eggs
- After a last vaccine planned 1 week before the transfer
- After de-worming of the flock (3 days prior the transfer)

In order to minimize the stress at transfer time, it is important to:

- Rear the birds on a similar drinking system as they will encounter on transfer.
- Increase light intensity to encourage water consumption
- Maintain temperature as close as close as possible to the temperature experienced by the pullets at the end of the rearing period.

4.3. LIGHTING PROGRAMME DURING THE PRODUCTION PERIOD

The light duration after transfer should be adjusted to match the light duration experienced at the end of the rearing period. The post transfer light duration should be at least the same length as during the rearing phase. It may be longer according to the plan for light stimulation.

As the birds remain sensitive to changes in light duration, never decrease the day length during the entire production period.



4.4. LIGHT INTENSITY MANAGEMENT DURING THE PRODUCTION PERIOD

It is possible after the peak of lay to reduce progressively the artificial light intensity. This may limit feed wastage, excessive activity of the birds and reduce the risk of mortality. Please take into account that light intensity should remain well distributed all over the house.

4.5. LIGHT TRAINING STRATEGY

In production several systems and lighting equipment are used. In all cases, target is to progressively switch off the light in the system and house to ensure birds move up to the perches areas.

Herbelow is an example of lighting strategy based on 3 lighting steps:

- Step 1: Switch off the light inside the system. At this way the hens are stimulated to move from under the system to the outside or above.
- Step 2: Switch off the side light outside the system
- Stpe 3: The central light on top of the system can be dimmed a few minutes minutes later.

Total procedure should take approximately 15 minutes, timing must be adapted according to birds behaviour.

Figure: Example of 3 steps light dimming procedures in production





4.6. FLOOR EGGS PREVENTION

Floor eggs is a multifactorial problem often related to management issue at the onset of lay or house and equipment design. We have indicated herebelow some important recommandations to maintain a good nesting behaviour:

- At the start of production, it is very important to collect floor laid eggs frequently. We recommend collecting every hour until the early afternoon. Floor eggs that are not removed become an incentive for other hens to lay in the same place. During this period, the farm attendant should try to identify those hens laying on the floor and place them onto the nests.
- Hens should not be disturbed while laying: avoid flock inspection and feed distribution during the main laying period
- Nest:
 - o Sufficent numbers
 - o Easy access
 - o Comfortable floor: use material such as Astro Turf, rubber floor, litter...
 - o Uniform repartition within the house
 - o Entrance of the nest should be well lighted and the inside of the nest darker
 - o No draught in the nest area
 - Possibility to use a light nest to switch on 1 hour before the house light switch on. It enables the brids which are laying earlier to find the nest.
- Lighting: ensure a uniform and sufficient light in the whole house. Any darker area is favourable to a floor laying behaviour.
- Birds should sleep on top of the system and not on the litter
- Litter: depth at the onset of lay should not exceed 2-3 cm
- Fence: when allowed by local regulation, an electric fence around the outside of the litter or slatted areas and along the walls and partitions may be useful. It should be fixed at 5 cm from the wall and 12 cm above the litter.
- Drinking and feeding systems in floor system:
 - o It should not be located more than 2-3 meters of the nest
 - Feeding system should be located as high or low as possible (according to the system), to allow an easy circulation of the hens. Chain feeders can be directly fixed on the slats.
- Specific case of Floor-Slats system:
 - o Stricly respect the proportion of 2/3 slats and 1/3 floor
 - o Possibility to use ladders to facilitate access to the slats
 - o Install the drinking and feeding system in the slats area
 - o After transfer, possibility to block the hens in the slats area for a maximum of 3 to 5 days. In this situation, be careful to the nest access.



4.7. MANAGEMENT OF EGG WEIGHT

The egg weight profile of a flock is mainly determined by the following factors:

- Bodyweight at light stimulation (or at sexual maturity).
 - The higher the bodyweight at sexual maturity the higher will be the egg weight during the laying period.
 - o The lower the bodyweight at sexual maturity the lower will be the egg weight during the laying period.
 - o Delay the onset of the egg production to increase average egg weight during the production period.
 - o Plan for an earlier sexual maturity to decrease average egg weight during the production period.
- Evolution of the bodyweight during the first week of production
- Nutrition also has important effects on the evolution of the egg weight during production:
 - o Intake of protein or digestible amino acids
 - o Metabolisable energy of the feed
 - o Linoleic acid and oil content of the feed

Graph: Cumulative NOVOgen Brown Light egg distribution at 90 weeks according to EU egg weight classification





4.8. MANAGEMENT OF EGG SHELL QUALITY

Egg shell quality has become even more critical due to the lengthening of production cycles and the development of automatic egg collection systems. While substantial genetic progress has been achieved, diet and feeding techniques are key to the expression of this genetic progress.

- Egg shell quality and colour depend essentially on the hen's ability to use its dietary calcium during shell formation i.e. shortly before night time until oviposition. A good calcium diet during the production period improves egg shell quality and colour and prevents bone demineralization (osteoporosis) and fractures. The quality of the calcium in the diet is determined by the calcium presentation, i.e. particle and / or powder form within the ration and its content in the feed. It is particularly important to increase the calcium content after 45 weeks to satisfy the increase of calcium requirement due to bigger egg size and to compensate for the natural shell quality decreasing with the age.
- Oyster shell is a good ingredient to improve the egg shell quality. The particle size and solubility are usually very good for laying hensit is possible to add 1g per bird at the end of the afternoon to promote a better calcium intake for supplying the egg shell deposition during the night. Take care of the quality about bacteriological level & calcium solubility to be sure to use a safe product.
- The feed distribution has also important effect on the shell quality. When possible, hens ingest more than 50% of their ration during the 6 hours prior to lights out in order to satisfy their specific calcium appetite. A fraction of this calcium intake is stored in the gizzard. Since hens do not eat during the night (calcification period), shell quality depends on the quantity of calcium remaining in the digestive tract at the end of the day. Another way to improve shell quality is to encourage hens to consum calcium in the middle of the night with the use of a flash feed when it is allowed. Please seek the advice of your local NOVOGEN technician.
- More information on the relation between feed and egg shell quality can be found in the NOVOGEN nutrition guide.

Eggshell quality (Newton)	30 - 40 weeks	40 - 50 weeks	50 - 60 weeks	60 – 70+ weeks
Good	> 41	> 40	> 39	> 38
Moderate	39	38	37	36
Poor	< 38	< 37	< 36	< 35

Table: Egg shell quality references per age

Photo: Shell strength measure





MONITORING BODYWEIGHT AND UNIFORMITY

5. MONITORING BODYWEIGHT AND UNIFORMITY

The main objective is to reach the appropriate bodyweight and uniformity targets at different stages of the bird's development:

- At the early stage (0 7 weeks: period of frame development)
- At sexual maturity with an even growth curve (a low bodyweight at sexual maturity could affect later performances)
- At the start of lay to the peak of production (a growth of at least 300g from 5% of lay until 30 weeks means that the bird's needs for egg production and growth are covered)

5.1. BODYWEIGHT CONTROL

- The birds must be sample weighed weekly from the first week. During the first 4 weeks, collective weights can be taken in batches of 5 or 10 birds using a bucket. Subsequently, the birds can be weighed individually.
- From 26 weeks of age, weigh the birds every 2 weeks and then monthly from 35 weeks of age,
- Weigh a sufficient number of birds (around 100) cornered using lightweight screens or frames in 2 or 3 places in the house. For an accurate interpretation of the result, it is important to weigh all the birds caught in the sample. Weights can be recorded on a weighing sheet which is available from our technicians. It is good practice to walk the house first to stir the birds up to allow for a more realistic sample to be penned.
- After weighing, average body weight and uniformity are calculated and immediately plotted on NOVOGEN growing curve chart. The analysis of the growing curve helps to accurately adjust the feed allowance (the quantities indicated in our feed section are only to be considered as a guideline) and when required, to take the appropriate steps to correct the uniformity.

5.2. UNIFORMITY CONTROL

- The uniformity target is set to ensure 80 % of the body weights are in a range between + 10 and -10 % of the flock mean body weight.
- The following factors play an important role in achieving and maintaining good uniformity:
 - o access to feed and water (see equipment standards)
 - o health status of the flock
 - o disease and parasitism
 - o quality of beak treatment
 - o temperature and ventilation



BEAK TREATMENT

6. BEAK TREATMENT

6.1. BEAK TREATMENT MONITORING

- Beak treatment is sometimes undertaken where either light intensity cannot be controlled due to the design of the house or when other kind of challenges chronically stress out the birds. The beak treatment procedure is performed to prevent feather pecking and cannibalism under these conditions.
- Beak treatment is a delicate operation and should only be carried-out by well-trained and experienced operators. Poor beak treatment can affect the ability of the birds to eat and drink correctly and leads to unevenness. Attention should be paid to local regulations regarding beak treatment and it is advisable to seek veterinary advice to ensure the procedures are being correctly applied.
- Two different methods can be used for beak treatment. The first one consists of an infrared treatment of the beak at day-old in the hatchery by using a specific machine. The second method can be practiced at 7-10 days by using a hot blade. With this second method and under some specific conditions, where permitted, a second beak treatment may be undertaken at 8-10 weeks.
- Before beak treatment with hot blades (in countries where this is permitted):
 - o check that the birds are healthy
 - o do not treat the beaks when the birds are reacting to vaccinations
 - o add vitamin K to the drinking water (to prevent haemorrhaging)
 - o check that the temperature of the blades is high enough to prevent haemorrhaging, but not too high which may risk chicks being burned.
- To limit the effect of beak treatment with hot blades on the feed consumption and water intake, it is
 important to increase the water level in the drinkers and the pressure in the pipes. Ensure that the depth
 of the feed in the feeders is correct.

As outlined above, in addition to the technical recommendations, any local code or regulation concerning animal welfare should be respected.

Photo: Infra-red beak treatment machine





Photo: Pullet with a treated beak



BEAK TREATMENT

6.2. FULLY BEAKED FLOCKS MANAGEMENT

Photo: Fully beaked pullet



In case of fully beaked flocks, different measures are necessary to prevent the consequences on selective feed intake and potential damage due to pecking:

- Strictly respect the feeding and drinking spaces and the stocking density standards before starting up. It will ensure a good uniformity of the flock and avoid risks of competition and fights between the birds.
- Keep the birds calm by avoiding any stress: equipment malfunction, visitors, any sudden changes in the management (light intensity, number of feed distributions, feed formulation...).
- Provide distractions for the birds to keep them busy: wood shaving pack, oyster shell, plastic strings... All the toys could help to maintain a good behavior. But layers get used to the toys very quickly, they need to be changed regularly, each week for example, otherwise, they stop playing with.
- Distribution of grit and grain on the floor can also enhance litter scratching.
- There is a strong relationship between feeding time and pecking, the shorter the feeding time the higher the risk of pecking. Also, the feed presentation is of high importance. If the particles are too big, it will reduce the feeding time and increase the risk of feather pecking and feed selection. If they are too fine it will induce poor appetite.
- Be on the alert for the absence of fluff or small feathers on the floor. It can mean that the birds are eating the feathers due to a nutrient deficiency in fibre source and it can easily turn into feather pecking. It can be confirmed by the presence of feather in the intestine. It is sometimes possible to reduce the level of feather pecking by increasing the fibre content of the feed, particularly insoluble fibres. It is also possible to add alfalfa or straw on the floor to provide additional fibre for the bird.
- Feed has to be well balanced in energy, amino acid, nutrient, fibre... Any deficiency in the feed can induce feather pecking.
- Placing some gas concrete blocks in the house as they can help to smoothen the sharp end of the beak and thereby prevent pecking and selective feed intake.
- It is important to maintain a suitable climate (ventilation, temperature). Hens that feel uncomfortable are inclined to start feather pecking.
- Special attention has to be given to the lighting. Light intensity and duration are important for preventing
 pecking. It is sometimes possible to reduce pecking by dimming or colouring the lights. In dark house, it
 is important to avoid any unwanted light source from outside during the night period.
- In production, it is recommended to dim the light a few days after the adaptation in the production house and to keep the nest in a dark area.
- Respect the health program and avoid parasitic proliferation.



WATER MANAGEMENT

7. WATER MANAGEMENT

Water is the first and most important requirement for poultry, it drinks around 1.8 times more than it eats. Therefore, water management and quality is a priority to ensure optimal flock performances.

7.1. WATER MANAGEMENT DURING THE REARING PERIOD

Table: Water management recommendations

Period	Advice
	Check the quality of the drinking water and the pipelines.
Before start up	Clean and rinse the water pipelines.
	• Make sure that the water temperature is around 20 to 25°C.
	Check the nipples.
	Provide unlimited water.
Δfter start un	• Keep the water pressure as low as possible. It will help the chicks to find the water
	• Adjust the height of the drinkers according to the chick size and growth. The chicks must be able to drink without difficulties.

7.2. WATER QUANTITY

The ratio water/feed is generally mentioned as being close to 2.0, but in reality it depends more on the environmental temperature. In a hot temperature climate, this ratio increases as the birds will drink more and eat less. In this condition, it is recommended to supply cool water to the birds. Water temperature above 20 °C should be avoided.

Water consumption is an important indicator to follow. Therefore, it is recommended to install a water meter on the water line. This equipment is inexpensive and easy to install. A low consumption can indicate a shortage of water supply or a sanitary problem on the flock. Waste of water can also be detected thanks to the water meter.

The below table gives some indication on water consumption according to the environmental temperature (source: ITAVI, 2012).

Table: Water consumption recommendations according to environmental temperature

House T ⁰C	Water/Fee	Water consumption in	
	Rearing	Production	production (ml/bird)
15	1.6	1.7	210
20	1.7	1.8	205
25	2.3	2.1	230
30	3.0	3.1	320

7.3. WATER QUALITY

- A water sample for analysis should be taken at the entry point of the house to check the quality of the water supply, and at the end of the system to check the efficiency of the disinfection system.
- Sample once or twice a year. More samplings should be performed especially when using a local water supply (surface well, deep well, etc...)
- Clean the pipe system during the sanitary break between flocks.

Clean drinkers on a regular basis

			WATER MANAGEMENT		
	Table	: Water quality meas	sures interpretation (ITAVI, 2007)		
	Crit	eria	Risk		
			Antibiotic and vaccine loss of efficiency		
		8 < Ha	Reduction of chlorine treatment efficiency		
	pH		Favours the growth of detrimental Gram negative bacteria (Salmonella, E. coli, etc)		
	(3.3 < pri < 0.3)		Urinary and/or digestive problems		
ochemical recommendations		pH < 4	Skeletal weakening		
			Water system corrosion		
			Reduces solubility of some antibiotics and vitamins		
	Hardness	> 20°F	Tartar development in the water system, but poultry is quite tolerant to high hardness		
	(10 to 15°F)	< 6°F	Trace minerals deficiency and influence on egg shell strength		
			Water system corrosion		
	Iron		Decrease in water intake		
	(≤ 0.2 mg/l)	Iron > 1 mg/l	Reduction in chlorine treatment efficiency		
ysic	Manganese	> 0.15 mg/l	Increases bacterial development		
Ph	(≤ 0.05 mg/l)				
	Nitrates	> 50 ma/l	Digestive troubles at high concentration		
	(≤ 50 mg/l)	00g.	Reduces vaccine efficiency		
	Organic matter	> 5 ma/l	Look for source of contamination (water supply, biofilm)		
	(≤ 2 mg/l)	e night	(
	Nitrites	> 0.1 mg/l	Enhances biofilm development		
	(≤ 0.1 mg/l)		Can be toxic at low concentration		
S	Total flora	> 100 germs/ml			
ical tion	(≤ 100 germs/ml)	gennenni	Faecal germs contamination can be detrimental in itself as		
olog nda	Salmonella	> 0 germs/ml	a direct source of pathogenic agents (ex: E. coli), but also		
teric nme	(0 germs/ml)	> 0 germa/mi	serve as an indicator for other contaminants (like parasites		
Bac econ	E. Coli	> () germs/ml			
۲.	(0 germs/ml)	5 gornio///ii			

Before starting any water treatment, it is recommended to take advice from a local specialist.



WATER MANAGEMENT

Table: Main physicochemical treatments

	рН	Hard	ness	Iron / Manganese
Treatment	Mineral acidification: chloride or sulphuric acid Organic acidification: formic, propionic, lactic acid	Neutralization (water < 10°F)	Softening (water > 15°F)	Iron / Manganese removal
Effect	pH reduction Small dose can be enough Bacteriostatic or bactericidal action depending on the acid used Potentially positive effect on digestive system (not for the mineral acids)	Increase of pH and hardness	Hardness reduction	Iron / Manganese removal

Table: Main antibacterial treatments

	Chlorine	Chlorine dioxide	Hydrogen peroxide
Principle of action	Mixed in the water Free residual chlorine (FRC) is reacting with bacteria and has to be measured at the end of the water line.	Soluble gas made from the mix of sodium chlorite and chlorehydric acid. The gas is injected in the water.	Mixed in the water Recommended to be used with paracetic acid for the pipes cleaning during the flock (intermittent treatment)
Methods	Initial dosage of 1 to 4 mg/l of active chlorine 0.3 to 0.6 mg/l of free chlorine at the end of the pipe line. To check every 2 weeks with a DPD coloured system. For a better efficiency, avoid high level of iron, manganese and organic matter and a pH > 7.5.	Maximum initial dosage of 1 mg/l 0.3 to 0.5 mg/l of FRC at the end of the pipe line. To check every 2 weeks (reactant DPD). For a better efficiency, avoid high level of iron, manganese and organic matter. No effect of the pH on the treatment	Initial dosage depends on the product concentration 30 to 50 mg/l of hydrogen peroxide at the end of the water line. For a better efficiency, avoid the presence of organic matter. No effect of the pH on the treatment
Installation and running cost		++	++
Monitoring the efficiency	At the end of the water line with a colorimetric test: reactant DPD	With a colorimetric test: reactant DPD (level of chlorine dioxide = 1.9 free chlorine)	Level of hydrogen peroxide to be measured with reactive strips.
Efficient against	Bacteria (at a concentration of 0.2 mg/l at the end of the water line), fungi, algae and virus.	Bacteria, fungi, algae, virus and spores.	Bacteria, fungi, algae, virus and spores.
Other existing	water disinfection systems: electr	olysis, UV sterilization, Copper su	Iphate

8. NUTRITION

Some important points :

- All NOVOGEN recommendations are made with a quality protein. That means with a good availability and digestibility of Amino Acids (A.A.). Any quality degradation (see raw materials chapter) must result in a proportional decrease in the A.A. matrix value in order to avoid any deficiency that would limit the genetic potential of the animals.
- Any anti-nutritional factors that may alter the bioavailability of A.A. must be controlled, taken into account in terms of risk and managed accordingly.
- There are many methods of defining available phosphorus, in particular because of the basic reference which may be different (monocalcium, anhydrous dicalcium, monosodium...). It is therefore necessary to adjust locally according to the field results.
- Chlorine level is more important to follow in alternative systems (you should be more strict). The maximum recommendation is in case of good droppings quality. If the droppings are too wet, you must decrease this maximum (in some case, until the minimum of sodium).
- These recommendations are given for an average temperature in husbandry between 20 and 22°C, an optimal breeding and material. That means when the birds will go outside (with lower temperature), they need more energy compare to other system of production. It is recommended in this case to increase the energy level with at least 1.8% more of energy (should be adjusted according to the situation).
- It is recommended, whenever possible, to formulate feeds in A.A-ratios, on the basis of lysine in order to avoid any imbalance (causing a greater need and losses due mainly to competition between A.A.).

	Pullets	Laying hens
Dig. Lysine	100	100
Dig. Methionine	48 / 50	52 / 58
Dig. Methionine+Cystine	76 / 80	85 / 90
Dig. Tryptophan	19 / 20	20 / 22
Dig. Threonine	68 / 74	66 / 74

These ratios are indicated by ranges, they are adjusted according to the observed field performances and the desired production objectives (typically feather, egg weight, etc.).

- Attention to the cumulative potential enzymatic valuations on A.A., protein and energy (see chapter enzymes in NOVOGEN nutrition guide).
- In comparison to cage system, it is recommended to increase energy level by 50 Kcal, protein by 0.5% and crude fiber by at least 1% at the onset of lay.

A minimum total fiber level is recommended for all their positive effects on the digestion and behaviour of the animals. The minimum level at 4.5% should be increase according to birds behavior. Increase at least to 0.5% or 1% if you begin to see any signs of aggression between birds.



8.1.	EXAMPLE OF	DIET SPECIFI	CATIONS FOR	REARING PERIOD
------	------------	--------------	-------------	-----------------------

	Starter	Grower	Pullet / Developer	Pre-Lay
	0 - 5 weeks	6 - 10 weeks	11 - 15 weeks	16 weeks to 2%
	Crumble	Crumble or Mash	Coarse Mash	Coarse Mash
EM Kcal / Kg	2900 - 3000	2800 - 2900	2700 - 2800 (1)	2700 - 2750 (1)
EM Kcal / lb	1316 - 1362	1270 - 1316	1225 - 1270	1225 - 1247
% Crude protein	20.0 - 21.0	18.0 - 19.0	16.0 - 17.0	16.0 - 17.0
% Crude fiber	-	2.5 - 5.0	5.0 - 6.5	4.0 - 5.5
% Crude fat	3.5 - 5.5	3.0 - 4.5	2.5 - 4.0	2.5 - 4.5
% Tot. Lysine	1.12	0.97	0.75	0.81
% Tot. Methionine	0.51	0.45	0.35	0.41
% Tot. Methio + Cystine	0.86	0.76	0.67	0.70
% Tot. Tryptophan	0.22	0.20	0.17	0.19
% Tot. Threonine	0.77	0.67	0.57	0.59
% Dig. Lysine	1.00	0.86	0.67	0.72
% Dig. Methionine	0.48	0.42	0.33	0.37
% Dig. Meth & Cystine	0.76	0.67	0.58	0.62
% Dig. Tryptophan	0.19	0.17	0.15	0.16
% Dig. Threonine	0.68	0.58	0.48	0.50
% Calcium	1.00 - 1.10	1.00 - 1.10	0.95 - 1.05	2.20 - 2.50
% Available Phosphorus	0.45 - 0.50	0.40 - 0.45	0.37 - 0.40	0.42 - 0.45
% Sodium	0.17 - 0.20	0.16 - 0.18	0.16 - 0.18	0.17 - 0.19
% Chlorine	0.16 - 0.20	0.16 - 0.20	0.16 - 0.20	0.16 - 0.22
% Potassium	0.70 - 0.80	0.65 - 0.80	0.60 - 0.80	0.60 - 0.80

(1) The energy level of the pullet feed 11 - 15 weeks and pre-lay should be equal to that of the laying feed at the beginning of lay.

(2) In hot climates, it is recommended to increase amino acid levels by 5% in order to compensate for lower consumption.

(3) If the birds go outside, energy level should be a little bit higher (must be adjusted according to the external temperature)



8.2. EXAMPLE OF DIET SPECIFICATIONS FROM 2% PRODUCTION TO 28 WEEKS

	Layer 1					
Ingested quantity (g/d)	Need g/ bird / day	100 (1)	105 (1)	110 <i>(1)</i>	115 <i>(1)</i>	120 (1)
% Crude protein	20.0	20.0	19.1	18.3	17.5	16.8
% Crude fiber	-			4.5 - 6.0		
% Crude fat	-			2.5 - 5.5		
% Tot. Lysine	-	0.96	0.91	0.87	0.83	0.80
% Tot. Methionine	-	0.49	0.47	0.45	0.43	0.41
% Tot. Methio + Cystine	-	0.82	0.78	0.75	0.71	0.68
% Tot. Tryptophan	-	0.22	0.21	0.20	0.19	0.18
% Tot. Threonine	-	0.69	0.66	0.63	0.60	0.58
% Tot. Isoleucine	-	0.85	0.81	0.77	0.74	0.71
% Tot. Valine	-	0.90	0.86	0.82	0.78	0.75
% Dig. Lysine	0.85	0.85	0.81	0.77	0.74	0.71
% Dig. Methionine	0.46	0.46	0.44	0.42	0.40	0.38
% Dig. Meth & Cystine	0.73	0.74	0.70	0.67	0.64	0.61
% Dig. Tryptophan	0.19	0.19	0.18	0.17	0.16	0.16
% Dig. Threonine	0.59	0.59	0.56	0.54	0.51	0.49
% Dig. Isoleucine	0.77	0.77	0.73	0.70	0.67	0.64
% Dig. Valine	0.82	0.82	0.78	0.74	0.71	0.68
% Calcium	4.20	4.20	4.00	3.80	3.65	3.50
% Available Phosphorus	0.42	0.42	0.40	0.38	0.37	0.35
% Sodium Min.	0.17	0.17	0.17	0.17	0.17	0.17
% Chlorine Max.	0.22	0.22	0.22	0.22	0.21	0.20
% Linoleic acid (Min.)	-	1.30	1.25	1.20	1.15	1.10

(1) The consumption levels listed above correspond to the usual observed consumption after 25 weeks.

For this period, we recommend to increase by 7% protein and amino acids to take into account both the needs of growth in addition to production needs and secondly, to take account of lower consumption observed during the beginning of this period.



8.3. EXAMPLE OF DIET SPECIFICATIONS FROM 28 TO 45 WEEKS

	Layer 2					
Ingested quantity (g/d)	Need g/ bird / day	100	105	110	115	120
% Crude protein	19.0	19.0	18.1	17.3	16.6	15.9
% Crude fiber	-		I	4.5 – 7.0		
% Crude fat	-			2.0 - 4.5		
% Tot. Lysine	-	0.90	0.86	0.82	0.78	0.75
% Tot. Methionine	-	0.46	0.44	0.42	0.40	0.38
% Tot. Methio + Cystine	-	0.77	0.73	0.70	0.67	0.64
% Tot. Tryptophan	-	0.21	0.20	0.19	0.18	0.18
% Tot. Threonine	-	0.64	0.61	0.58	0.56	0.53
% Tot. Isoleucine	-	0.79	0.75	0.72	0.69	0.66
% Tot. Valine	-	0.85	0.81	0.77	0.74	0.71
			I			
% Dig. Lysine	0.80	0.80	0.76	0.73	0.70	0.67
% Dig. Methionine	0.43	0.43	0.41	0.39	0.37	0.36
% Dig. Meth & Cystine	0.69	0.69	0.66	0.63	0.60	0.58
% Dig. Tryptophan	0.18	0.18	0.17	0.16	0.16	0.15
% Dig. Threonine	0.55	0.55	0.52	0.50	0.48	0.46
% Dig. Isoleucine	0.72	0.72	0.69	0.66	0.63	0.60
% Dig. Valine	0.77	0.77	0.73	0.70	0.67	0.64
			I			
% Calcium	4.20	4.20	4.00	3.80	3.65	3.50
% Available Phosphorus	0.40	0.40	0.38	0.36	0.35	0.33
% Sodium Min.	0.17	0.17	0.17	0.17	0.17	0.17
% Chlorine Max.	0.22	0.22	0.22	0.22	0.22	0.22
% Linoleic acid (Min.)	-	1.30	1.25	1.20	1.15	1.10

Fiber level must be adjusted according to the bird behavior (cf. our Novogen nutrition guide for more details)



8.4. EXAMPLE OF DIET SPECIFICATIONS FROM 45 TO 70 WEEKS

	Layer 3					
Ingested quantity (g/d)	Need g/ bird / day	100	105	110	115	120
% Crude protein	18.5	18.5	17.7	16.9	16.2	15.5
% Crude fiber	-		1	4.5 - 7.5		
% Crude fat	-			1.5 - 3.5		
% Tot. Lysine	-	0.90	0.86	0.82	0.78	0.75
% Tot. Methionine	-	0.46	0.44	0.42	0.40	0.38
% Tot. Methio + Cystine	-	0.77	0.73	0.70	0.67	0.64
% Tot. Tryptophan	-	0.21	0.20	0.19	0.18	0.18
% Tot. Threonine	-	0.64	0.61	0.58	0.56	0.53
% Tot. Isoleucine	-	0.79	0.75	0.72	0.69	0.66
% Tot. Valine	-	0.85	0.81	0.77	0.74	0.71
% Dig. Lysine	0.80	0.80	0.76	0.73	0.70	0.67
% Dig. Methionine	0.43	0.43	0.41	0.39	0.37	0.36
% Dig. Meth & Cystine	0.69	0.69	0.66	0.63	0.60	0.58
% Dig. Tryptophan	0.18	0.18	0.17	0.16	0.16	0.15
% Dig. Threonine	0.55	0.55	0.52	0.50	0.48	0.46
% Dig. Isoleucine	0.72	0.72	0.69	0.66	0.63	0.60
% Dig. Valine	0.77	0.77	0.73	0.70	0.67	0.64
			I			
% Calcium	4.50	4.50	4.30	4.10	3.90	3.75
% Available Phosphorus	0.38	0.38	0.36	0.34	0.33	0.32
% Sodium Min.	0.17	0.17	0.17	0.17	0.17	0.17
% Chlorine Max.	0.22	0.22	0.22	0.22	0.22	0.22
% Linoleic acid (Min.)	-	1.20	1.15	1.10	1.05	1.00

Fiber level must be adjusted according to the bird behavior (cf. our Novogen nutrition guide for more details)



8.5. EXAMPLE OF DIET SPECIFICATIONS FROM 70 WEEKS TO DEPLETION

	Layer 4					
Ingested quantity (g/d)	Need g/ bird / day	100	105	110	115	120
% Crude protein	18.0	18.0	17.2	16.4	15.7	15.1
% Crude fiber	-			4.5 - 7.5		
% Crude fat	-			1.5 - 3.0		
% Tot. Lysine	-	0.90	0.86	0.82	0.78	0.75
% Tot. Methionine	-	0.46	0.44	0.42	0.40	0.38
% Tot. Methio + Cystine	-	0.77	0.73	0.70	0.67	0.64
% Tot. Tryptophan	-	0.21	0.20	0.19	0.18	0.18
% Tot. Threonine	-	0.64	0.61	0.58	0.56	0.53
% Tot. Isoleucine	-	0.79	0.75	0.72	0.69	0.66
% Tot. Valine	-	0.85	0.81	0.77	0.74	0.71
% Dig. Lysine	0.80	0.80	0.76	0.73	0.70	0.67
% Dig. Methionine	0.43	0.43	0.41	0.39	0.37	0.36
% Dig. Meth & Cystine	0.69	0.69	0.66	0.63	0.60	0.58
% Dig. Tryptophan	0.18	0.18	0.17	0.16	0.16	0.15
% Dig. Threonine	0.55	0.55	0.52	0.50	0.48	0.46
% Dig. Isoleucine	0.72	0.72	0.69	0.66	0.63	0.60
% Dig. Valine	0.77	0.77	0.73	0.70	0.67	0.64
% Calcium	4.80	4.80	4.60	4.40	4.20	4.00
% Available Phosphorus	0.36	0.36	0.34	0.33	0.31	0.30
% Sodium Min.	0.17	0.17	0.17	0.17	0.17	0.17
% Chlorine Max.	0.22	0.22	0.22	0.22	0.22	0.22
% Linoleic acid (Min.)	-	1.20	1.15	1.10	1.05	1.00

Fiber level must be adjusted according to the bird behavior (cf. our Novogen nutrition guide for more details)



9. HEALTH PROGRAMME

It is impossible to devise a health programme to adequately suit all geographic areas. For this reason, it is strongly recommended to consult a local specialist to establish a prevention programme adapted to the region.

This guide limits its comments to the description of some rules for the use of vaccines and other treatments. To be successful, respecting these rules is as important as choosing the right products:

- Staff should be properly trained to carry out veterinary operations. It is useful to create a Standard Operating Procedure Manual, that describes in full details the way to perform each vaccination or treatment.
- All the necessary equipment (sprayers, syringes, etc.) must be correctly maintained, and checked before each use.
- Each operation should be planned and supervised by a technically competent person.
- Vaccines and treatments should be stored in appropriate conditions, in suitable quantities considering the requirements and supply time.
- Report carefully in the flock records the details of all operations: date, time, vaccine batch number, route, etc.
- Finally, it is useful to have the help of a laboratory in order to anticipate health problems ahead of time and to assess the efficiency of the :
 - o control of disinfection, water and feed quality
 - o serological monitoring
 - o post mortem examination, routine parasite checks

9.1. VACCINATION

The vaccination programme has to be defined and regularly updated with a local veterinarian.

- Vaccine rules:
 - o Only vaccinate healthy flock
 - o Always check the conformity of the vaccines with the vaccination programme. In case of doubt, immediately contact your local veterinarian.
 - o Keep records of vaccine serial numbers and expiry dates
 - o Use clean and specific vaccination equipment
- Vaccine preparation:

Live vaccine	Inactivated vaccine
 Live vaccines are fragile and should be prepared with care. Vaccine storage should be monitored, to make sure it is kept as per manufacturer recommendations (usually between 2 to 6°C). Regularly check the temperature of the storage equipment. For freeze-dried vaccines, dilute the powder into mineral water or with the provided diluent if appropriate carefully injected with a syringe. The quality of the water used for live vaccine should be checked: no disinfectant, suitable amount of iron, manganese and magnesium, no organic matter. 	 Inactivated vaccines are administered through adjuvants. The adjuvant can shock the birds if injected too cold, so it is advised to slowly warm them up before use (25 to 30°C). To ensure quality injection, needles should be sterile before use, and changed regularly. A good base number is to change every 1,000 birds, but more importantly make sure that it is not damaging the muscle as it would cause unnecessary pain and be detrimental to the vaccine up- take.



	Tables: Vaccination methods and description		
VACCINATION METHODS	GENERAL DESCRIPTION		
	• Do not use disinfectant or chlorine in a period of 48 hours before vaccination and 24 hours after.		
	 Check each drinkers condition and cleanliness 		
	• Cut the water around 2 hours before vaccination. It also depends on the current temperature (one should be more cautious in hot climates and use lesser time)	Photo : Drinking water vaccination installation	
	Prepare the vaccine:		
	- Clean your hands		
Drinking water	- Prepare the required volume of water in a clean tank. A good base number is 15 to 20% of the previous day water consumption.		
	- Neutralize the residual chlorine with Sodium Thiosulfate (16g/1000l water) mix and let it act for 10 minutes.		
	- Mix the solution vaccine + diluent with the neutralized water for 1 minute	A. F.M	
	Distribute the vaccine	Photo : Coloured	
	• Let the birds drink the vaccine solution. It has to be consumed within 2 hours	tongue	
	• During the distribution, walk among the birds to move them and stimulate drinking behaviour.		
	• Once the vaccination solution has been consumed, give water without chlorine		
	• To check the proper vaccine take, a water colorant can be used. More than 90% of the birds should have a coloured tongue after the distribution.		
	• It is recommended tospray in the morning and in a calm atmosphere (avoid feed distribution right after the vaccination and dim the light).	Photo : Spray vaccination	
	Gather the birds		
	• Turn off the ventilation and heating systems to avoid losses due to evaporation or dispersion		
	Preferably use mineral water		
Spray	Check the condition and cleanliness of the spraying machine		
	Prepare the vaccine and fill up the spraying machine tank with the solution of vaccine and water		
	• Spray at 30-40 cm high / Make sure the droplets size is adapted to the vaccine used / Spray along the whole building length back and forth/ Do not spray if the birds pills up	A synthesis	
	• Wait for 5-10 minutes before switching the light, heating and ventilation systems back on		



HEALTH PROGRAMME			
VACCINATION METHODS	GENERAL DESCRIPTION		
	 Use specific vaccination equipment 		
	 Set the syringe at the required dose: the dose corresponds to the dose of vaccine to be injected or, for mixed vaccine, the sum of the doses of mixed vaccines 		
Subcutaneous	Hold the birds by the wings and inject the vaccine at the indicated location		
/ intramuscular injection	 Regularly check the consumed volume of vaccine according to the number of vaccinated birds 		
	 Beware of defusing syringe when the bottle is getting empty 		
	Regularly check the needle condition		
	Comfortable working conditions are the key to a successful vaccination		
		Photo : Eye drop vaccination	
	Prepare the vaccine		
Eve drop	 Hold the bird to be vaccinated with the head tilted to one side 		
Eye diop	 Drop one drop of vaccine into the eye. 		
	• Be sure the vaccine spreads over the eye before releasing the bird.		
	Use the provided stylet		
	Hold the bird on the side and spread the wing		
Wing web	• The ideal transfixion area is in the wing membrane facing the elbow		
-	• Dip the stylet in the vial and insert it through the wing.		
	Avoid damaging the blood vessels.		



Tables: Applications of vaccines

(indicative only, check with your local veterinarian)

Basic vaccines applications			
Disease	Administration methods	Application periods	
Marek	Intramuscular / subcutaneous / in-ovo	Day-old	
Newcastle Disease (ND)	Drinking water / Spray / Subcutaneous / Intramuscular / in-ovo	Depending on the local epidemiological context this can start at day 1	
Gumboro Drinking water / in-ovo		Depending on the local epidemiological context and/or quantity of antibodies of maternal origin	
Infectious Bronchitis (IB)	Drinking water / Spray / Subcutaneous / Intramuscular	Depending on the local epidemiological context, usually at day 1 with regular boosters	
Avian encephalomyelitis (AE)	Drinking Water / Wing Web	Usually around 12 to 14 weeks of age	

Optional vaccines applications			
Disease	Administration methods	Application periods	
Coccidiosis	Spray / drinking water	Day-old	
Infectious Laryngotracheitis (ILT)	Eye drop / Spray / Injection (recombinant vaccines) / in- ovo / Wing web	Depending on the vaccine and the local epidemiological context	
Fowl Pox	Wing Web	8 to 12 weeks	
Mycoplasmosis	Spray / Eye drop / Subcutaneous / Intramuscular	Depend on local epidemiological context and vaccine used	
Salmonella	Drinking Water / Spray / Intramuscular	Usually based on live vaccines 6 weeks apart and a booster with inactivated 4 weeks before lay	
Pasteurellosis	Wing web /Subcutaneous / Intramuscular	Depending on local epidemiological context	
Infectious Coryza	Subcutaneous / intramuscular	Depending on local epidemiological context	
Egg Drop Syndrome (EDS)	Subcutaneous / Intramuscular	Usually inactivated vaccine before lay	



9.2. PARASITES AND INSECTS MONITORING

The below tables give some indications on the main parasites and insects source of troubles in a layer poultry farm. The treatments have to be defined and regularly updated with a local veterinarian.

Table: Main parasites and insects in layer poultry farm

		DESCRIPTIONS	SIGNS	TREATMENT
	Ascaridia galli	 Roundworm that can measure up to 12 cm It is the most common layer paraite Adult female lays eggs in the intestine that pass in the faeces. Contamination by eggs ingestion, the larvae reaches the infectious stage after 2-3 weeks. The adult can live for one year. 	- anaemia, intermittent diarrhoea, bodyweight loss, egg production decrease, loss of fertility in males	 Litter sanitation measures Strict cleaning and disinfection between two flocks Chemical treatment: Benzimidazoles, avermectins, levamisole, etc
Internal parasites	Capilaria	 Threadworm that can measure up to 8 cm Parasite from the digestive tract located in the oesophagus, crop, small intestine or caecum according to the species. The eggs pass in the faeces and reach the infectious stage within 3-4 weeks After ingestion, it can produce sever inflammation and sometimes haemorrhage More common problem in deep litter houses 	 Young birds are more sensitive Bodyweight loss, apathy, egg production decrease, can lead to death 	 Litter sanitation measures Strict cleaning and disinfection between two flocks Chemical treatment: Benzimidazoles, avermectins, levamisole, etc
	Cestodes	 Tapeworm that can measure up to 4 cm. Their development cycle includes an intermediate host (insect, snails, slugs, beetles, ants, earthworms, houseflies). The hens are being infected by eating this host. Consequently, this parasite is uncommon in closed house farm. Once in the intestine, it reaches its maturity at around 3 weeks of age. 	- In case of heavy infection: bodyweight loss, feed intake decrease specially with young birds, egg production decrease	 Litter sanitation measures Strict cleaning and disinfection between two flocks Chemical treatment is usually based on praziquantel



		DESCRIPTIONS	SIGNS	TREATMENT
				- Chemical products:
External parasites	Red Mites	 Blood sucking during night periods Remain hidden in cracks and crevices during the light periods. About a day after feeding, the female lays eggs in cracks and crevices of the house Fast increase of population 	 Behaviour modification due to the disturbance (pecking, nervousness) Egg production decrease Anaemia that can induce higher mortality and FCR increase Blood stains on the egg shell To be detected and treated as soon as possible to avoid heavy infection by using specific trap 	Organophosphorus based (phoxim, azametiphos, dichlorvos) Pyrethroids based (cyfluthrin, permethrin) Spinosad based - Natural products: Silica, sodium bicarbonate, extracts and essential oils of medicinal and aromatic plants - Cyclic lighting programme (when allowed) - Importance of cleaning and disinfection measures between
Insects	Fly (Musca domestica)	 Female can lay up to 1000 eggs and start laying 4 to 8 days after mating. Egg development is optimum in manure with 40-70% humidity. Larvae are feeding on decomposing organic matter Adults can live 2 weeks in summer and up to 2 to 3 months in winter. 	 Passive agent of pathogens spread (virus, bacteria, parasites) Disturb the hens and reduce the egg production Flies dropping increase the number of second grade eggs 	 Maintain less than 25% water in the manure (ventilation, avoid waste of water, proper broken eggs disposal, regular removal of the manure) Chemical treatment should target both adults (organophosphates, pyrethroids, carbamates) and larvae (cyromazine, triflumuron, some organophosphates) Biocontrol: acaras, beetles, natural predators Alternate the active molecules to limit risks of resistance
	Darkling eetles	 Female can lay up to 800 eggs Larvae burrow into the ground for insulation as they pupate Life cycle (from egg to adulthood) of 2 months to a year depending on season and temperature 	 Passive carrier of pathogens (Marek virus, salmonella, E. Coli, aspergillus) Penetrate in insulation equipment causing heavy damages 	- Strict cleaning and disinfection between two flocks - Chemical treatment should target adults (preferably on the walls), and the larvae (preferably on the litter, under the feeding pans and/or drinkers)





LITTER

10. LITTER

10.1. LITTER QUALITY

Depending on the system, litter can be used to cover the floor in rearing. The main materials used for litter are chopped straw (wheat, barley...) and wood shaving. It is recommended to use wood shaving coming from non treated wood. Depending on the local availabilities, other material like rice hull and cellulose pellet can also be used. The choice of the litter is an important compenent on the success of rearing by ensuring a confortable environment while avoiding potential diease conditions. Moreover, litter is used by birds for dust bathing, foraging and scratching, which help reduce feather picking.

Photo: Wood shaving litter in rearing



Different qualities must be taken into account for the choice of the litter:

- Thermal insulation from the cold coming from the ground. It depends of the litter material and depth. It is very important during the first week, as long as the chicks are poikilotherm.
- Abosrption of humidity from manure and birds breath. It is particulary important in cold and wet weather when ventilation is low.
- Low level of dust to avoid respiratory disease.
- Safe for the chicks by avoiding being a source of contamination (virus, bacteria, mould).
- Comfortable and not harmful for the chicks

Ensure the shed is fully warmed before distributing the litter. If the gap of temperature between the floor and the room is too important, ie the floor is not warm enough, litter will become stick and wet from below.

10.2. LITTER MANAGEMENT

In order to maintain an optimal litter quality all along the flock:

- Keep under control litter humidity
- Avoid any leaking or spillage from water system (nipples, bell drinkers...).
- Floor system:
 - o In rearing, litter depth should be 5 cm high
 - o Additional fresh litter can be distributed on top of the existing litter to keep low moisture content
- Aviary system:
 - o 2 cm of litter is sufficient.

o In production, excess of litter should be frequently removed to avoid the floor area to be too confortable as it could induce some floor laid eggs. It also allow to reduce dust content.



FINE DUST MANAGEMENT

11. FINE DUST MANAGEMENT

The presence of dust in production houses is an inescapable consequence of cage free production. The hens have more possibilities to move inside the house which increase physical interaction with their environment.

A research in the Netherlands from Winkel and Albert (2016) has showed that while layer market has been changing from 99% cage production in 1995, towards 95% alternative production in 2014 the fine dust emission (PM_{10}) by poultry farm in the country has been increasing by 5 times.

One of the main challenge for poutry farmers and the poultry sector is how to manage the fine dust production in order to avoid its impact on farm workers and public health. With the rise of alternative production and the increase of fine dust production in North-West Europe, climate and poultry equipment suppliers have specialized themselves in the search of fine dust reducing equipment. The available equipment can be categorised in 4 main groups, which are presented below. The investments and fixed annual cost are variating according to equipment.

Fine dust reducing equipment	Principal	Fine dust reducing capacity
Heat exchangers	The temperarure of the incoming air is increased with the in house temperature. Condensation will be produced at the heat exchanger which is mixed with fine dust. Other principals with dust filters are also available.	+
Filters	The inhouse air or the outgoing air goes over a filter which separate the dust from the air. Different filter systems (paper, biofilter, synthetic,etc.) are available. The effect differs, based on the air movement, the available filter surface and the frequency to change/clean filter systems	+++
Dry-or manure tunnel	Most alternative production houses have manure collecting belts at the back side of the production house. The manure must be dried before it can be transported for other purposes. The air from the production house with the fine dust can be used for this. Because the manure is wet, it will bind the fine dust.	++
lonization	The inhouse air or the outgoing air is ventilated over ionization equipement. The positive or negative tension creates a field of loaded iones which move to the grounded equipment. The fine dust binds to the loaded ionen.	+++

Table: Fine dust reducing equipment



PASTURE MANAGEMENT

12. PASTURE MANAGEMENT

Free range systems requires a specific pasture management in order to optimize its exploration by hens.

Benefits of a well-managed pasture are multiple:

- Animal welfare by allowing hens to express their natural exploration behaviour.
- Lower sanitary risk, specially parasite infections.
- Better distribution of faeces in the land reducing accumulation of phosphorus and nitrogen in crowded area.
- Lower hens physical effect on the grass.
- Optimization of landscape integration with plantations.

Pasture is not a natural environment for layers hens. Plantations and shelters are necessary to encourage hens to explore the pasture for the following reasons:

- Supplying of shadow area.
- Protection from the wind.
- Guide birds movement.
- Protection from rapacious.

12.1. OPTIMIZATION OF PASTURE EXPLORATION

Different areas can be distinguished requiring a specific management:

- House and its surroundings:
 - House should be slightly over elevated from ground level to avoid possible water infiltration inside the house which could alterate litter quality.
 - o Use connected gutter and drain around the house to reduce water and mud accumulation.
 - o A concrete pavement in fromt of the pop holes helps to keep a clean area.
- House exit area (up to 15m from the house)
 - o This is the first area explorated by hens.
 - o It must be clean, conformatbale and protected from the wind.
 - o A too shaded area will not be attractive for birds.
 - o To avoid feces accumulation, it should only be a transit area.
 - Intermediary and background area (from 15 to 40m from the house)
 - o Once hens get out of the house, target is to encourage them to explore further and reach the intermediary area.
 - o In these areas, plantations should provide shelters from predators, protect from the wind and provide shadow areas. It is often recommended to mix bushes and trees plantations.
 - o These areas should also be drained to avoid water accumulation.



PASTURE MANAGEMENT

12.2. MANAGEMENT OF PARASITIC RISK

Pasture is a risk area regarding pathogenes and parasites contaminations. They are brought by hens feces as well as rodents and wild birds.

Coccidiosis is often the main parasite found but ascaris, capilaria, tenia can also be present.

Logically, the most used area are the most contaminated. Therefore, it is recomendend to avoid environments favourable to the development of these parasites:

- Avoid any wet area with a good draining of the pasture, house surrending and gutter connections.
- Good quality fences reduce entrance of rodents.
- The most crowded area, specially house surrounding, must be disinfected between 2 flocks with quick lime (400kg / 1000m2) or caustic soda (50 to 100kg / 1000m2).



GENERAL FARM RULES

13. GENERAL FARM RULES

Ideally, the best rule of management is to have one age and one breed per site to ensure the "all-in, all-out" principle is followed at all times.

The choice of the site for the farm, including the layout of the houses, must prioritise the elimination of all possible sources of contamination. Biosecurity protection is reinforced by hygiene controls.



A changing room should be made available at the entrance of the site. It must be used by everybody entering the farm (incorporating both a shower and a change of clothes).



When the old flock is removed and before the arrival of the new flock, all houses and equipment must be thoroughly cleaned and disinfected according to strict procedures and protocols. This should be followed by a rest period of at least 10 days.



CLEANING AND DISINFECTION OF POULTRY HOUSES

14. CLEANING AND DISINFECTION OF POULTRY HOUSES

Between each flock, cleaning and disinfection of the houses, their annexes, surroundings and access ways are essential to ensure the optimal health conditions required for the incoming flock to maximise its profitability.

14.1. OPERATIONS PRIOR TO CLEANING

- Water tank, pipes and nipples:
 - Empty the complete water system.
 - Clean and de-scale the complete system with an acid solution and leave for 6 hours to soak.
 - Rinse twice with clean water.
- All the equipment (nests, feeders, drinkers etc.) are removed and stored on a concrete area.
- The entire ventilation system (air inlets and outlets, fans, heating and ventilation ducts if they are present) and individual radiant or pancake type brooders are brushed and vacuum cleaned.
- Litter is removed.

14.2. WASHING

When washing, ensure local regulations regarding wash down water are observed. As a rule, always ensure that the dirty water is directed towards a pit or suitable internal drain and does not run outside to the house surroundings or access roads and pathways.

✤ House

- Soak and remove the remaining organic matter
- Apply a bactericidal and fat removing detergent using an appliance capable of dealing with foam products.
- Some hours after soaking, wash with a high pressure washer (>50kg/cm²) or with hot water, in the following order:
 - o Internal roof surfaces, from the top downwards.
 - o Walls, from the top downwards.
 - o Finally, pits and concrete floors.
 - Equipment
- Nests, drinkers and feeding equipment:
 - o Soak and remove all organic matter.
 - o Apply a bactericidal and fat removing detergent using an appliance capable of dealing with foam products.
 - o Ensure every piece of equipment gets a thorough wash, followed by rinsing.
 - Prior to the final rinsing, immerse the removable parts of the nests (perches and nest box bottoms) for 24 hours in a disinfectant solution.
 - o Dry on a clean disinfected concrete area (different to that used for washing).



CLEANING AND DISINFECTION OF POULTRY HOUSES

14.3. PLACING EQUIPMENT BACK INTO THE HOUSE

The vehicles used for this operation must have been carefully washed and sprayed with disinfectant.

14.4. DISINFECTION

- Water pipes
 - o Prepare a highly concentrated chlorine solution (200 ppm) in the water tank.
 - Open the tank to fill the pipes with this solution and leave for 24 hours. Afterwards, drain the water circuit. Do not forget to seal the water tank to protect it from dust.
- House
 - House and equipment disinfection is achieved using a homologous bactericidal, virucidal and fungicidal disinfectant, applied with a hand held or low pressure sprayer or a foam-producing machine.
 - o The list of homologous approved disinfectants may vary from one country to another. We recommend that you consult the relevant local Authorities for a list of approved disinfectants and the required concentrations when used for poultry applications.
- Feed Storage Silos
 - Scrape, brush wash and after drying, fumigate using fungicidal candles following the manufacturer's guidelines.
- Heating and ventilation ducts (if they are present)
 - o Disinfection using fungicidal, virucidal and bactericidal candles following manufacturer's guidelines.
- House surroundings and road and path access ways.
 - o Spread a disinfecting product, such as:
 - caustic soda (50 to 100 kg/1000 m²).
 - or quicklime (400 kg/1000 m2).

14.5. SANITARY PRECAUTIONS

Place clean boots and overalls in the changing room. Replenish footbaths with an appropriate disinfectant.

14.6. ASSESSING DISINFECTION EFFECTIVENESS

- Visual examination
 - o Check for dirt stains in the house and on the equipment.
- Bacteriological analysis
 - Contact plates or swabs are applied to equipment and to different places in the house. These are rapidly forwarded to a laboratory for bacteriological assessment following an agreed protocol with the laboratory.

14.7. RODENT CONTROL

Rodents may be vectors of numerous bacterial diseases such as salmonellosis.

Rodent control is often based on the use of toxic baits which generally contain anticoagulants. These are left in places frequented by the rodents following a site risk assessment. A poorly prepared rodent control programme may give variable or poor results. We therefore advise using a specialised rodent control service.



Commercial layers – Alternative System Management Guide – NOVOgen BROWN Light

CLEANING AND DISINFECTION OF POULTRY HOUSES

14.8. RESTING PERIOD

This starts only when all the above operations have been achieved and lasts for at least 10 days, in order for the house to dry properly.

14.9. BEFORE THE NEW FLOCK ARRIVES

- 3 days before the new flock arrives, a residual insecticide is sprayed on all surfaces.
- Fresh litter is placed (never use mouldy material) and its surface sprayed with a larvicidal insecticide.
- Equipment is prepared in the brooding area.
- 24 hours before the new flock arrives, the final disinfection is performed by fogging.
- To be careful with disinfection fo vehicles coming to the farm





Caring for life NOVOGEN S.A.S – 5, RUE DES COMPAGNONS – SECTEUR DU VAU BALLIER – 22960 PLEDRAN - FRANCE

Tel. + 33 (0)2 96 58 12 60 - Fax + 33 (0)2 96 58 12 61

contact@novogen-layers.com

www.novogen-layers.com